Author Index

Aguilar, M. I., see Hodder, A. N. 17
Ahnoff, M.
——, Chen, S., Green, A. and Grundevik, I.
Chiral chloroformates as transparent reagents for the resolution of metoprolol enantiomers by reversed-phase liquid chromatography 593
Albert, K., see Pfleiderer, B. 343
Arvidsson, E.
——, Jansson, S. O. and Schill, G.
Chiral separations of atropine and homatropine on an α_1-acid glycoprotein-bonded stationary phase 579
Ascah, T. L.
—— and Feibush, B.
Novel, highly deactivated reversed-phase for basic compounds 357
Bartha, A.
——, Vigh, Gy. and Ståhlberg, J.
Extension of the electrostatic retention model of reversed-phase ion-pair chromatography to include the simultaneous effects of the organic modifier and the pairing ion 85
Bayer, E., see Pfleiderer, B. 343
Belliardo, F.
——, Chiantore, O., Berek, D., Novàk, I. and Lucarelli, C.
Development and use of carbon adsorbents in the liquid chromatographic separation of isomers 371
Berek, D., see Belliardo, F. 371
Bertucci, C.
——, Domenici, E., Uccello-Barretta, G. and Salvadori, P.
High-performance liquid chromatographic resolution of racemic 1,4-benzodiazepin-2-ones by means of a β-cyclodextrin silica bonded stationary phase 617
Boppana, V. K.
—— and Rhodes, G. R.
High-performance liquid chromatographic determination of guanidino compounds by automated pre-column fluorescence derivatization 279
Bristow, P.A.
Towards a chromatographic quantitator 265
Camacho, P. L.
——, Geiger, E., Vigh, Gy., Webster, R. and Thompson, D. H.
Separation of the enantiomeric intermediates of some platelet-activating factor analogues on naphthylalanine-type Pirkle column 611
Chen, S., see Ahnoff, M. 593
Cheung, A. P.
—— and Kenney, D.
Partition coefficients and capacity factors of some nucleoside analogues 119
Chiantore, O., see Belliardo, F. 371
Clark, B. J., see Marr, J. G. D. 289
Cserháti, T., see Valkó, K. 35
Czok, M.
—— and Guiochon, G.
Aligned fiber columns for size-exclusion chromatography 303
Davankov, V. A., see Kurganov, A. 391
Dohi, T., see Sakuma, I. 223
Domenici, E., see Bertucci, C. 617
Dominguez, E., see Marko-Varga, G. 423
Driessche, N. Vanden, see Maris, F. 211
Duijsters, P. P. E., see Tock, P. P. H. 185
Durand, A., see Rouchouse, A. 601
Edlund, P. O., see Johansen, K. 471
Eltekova, S. A.
Retention times and heats of adsorption of aromatic compounds on carbon adsorbents 335
Engelsma, M.
——, Kok, W. Th. and Smit, H. C.
Selective determination of trace levels of phenol in river water using electrochemical concentration modulation correlation chromatography 201
Eray, B., see Kurganov, A. 391
Eriksson, B.-M., see Gustafsson, S. 75
Farkas, Gy., see Vigh, Gy. 481
Feibush, B., see Ascah, T. L. 357
Fell, A. F., see Marr, J. G. D. 289
Fellegvári, I., see Valkó, K. 35
Fornstedt, T.
——, Westerlund, D. and Sokolowski, A.
Studies on system peaks in ion-pair adsorption chromatography. III. Regulation of system peak gradient retention for obtaining analyte peak compression 61
Frank, H., see Welsch, T. 97
Fujimoto, C., see Jinno, K. 443
Fukui, Y., see Sakuma, I. 223
Geiger, E., see Camacho, P. L. 611
Golshan-Shirazi, S.
—— and Guiochon, G.
Solutions of the equilibrium and semi-equilibrium models of chromatography 495
Gorton, L., see Marko-Varga, G. 423
Green, A., see Ahnoff, M. 593
Grundevik, I., see Ahnoff, M. 593

Guiochon, G., see Czok, M. 303
——, see Golshan-Shirazi, S. 495
Gustafsson, S.
——, Eriksson, B.-M. and Nilsson, I.
Multiple peak formation in reversed-phase liquid chromatography of ramipril and ramiprilate 75
Hann-Hägerdal, B., see Marko-Varga, G. 423
Hara, S., see Uchida, T. 327
Hayashida, M.
——, Nihira, M., Watanabe, T. and Jinno, K.
Application of a computer-assisted high-performance liquid chromatographic multi-wavelength ultraviolet detection system to simultaneous toxicological drug analyses 133
Hearn, M. T. W., see Hodder, A. N. 17
Hindriks, R., see Maris, F. 211
Hirata, N., see Noguchi, K. 145
Hodder, A. N.
——, Aguilar, M. I. and Hearn, M. T. W.
High-performance liquid chromatography of amino acids, peptides and proteins, XCVII. The influence of the gradient elution mode and displacer salt type on the retention properties of closely related protein variants separated by high-performance anion-exchange chromatography 17
Ishikawa, T., see Yamakawa, Y. 319
Izu, H., see Uchida, T. 327
Jalonen, H., see Örn, G. 627
Janssen, H.-G., see Schoenmakers, P. J. 563
Jansson, S. O., see Arvidsson, E. 579
Jinno, K.
—— and Fujimoto, C.
Coupling of microcolumn high-performance liquid chromatography with Fourier transform infrared spectrometry 443
——, see Hayashida, M. 133
Johansen, K.
—— and Edlund, P. O.
Determination of water-soluble vitamins in blood and plasma by coupled-column liquid chromatography 471
Kaliszan, R.
—— and Ośmiałowski, K.
Correlation between chemical structure of non-congeneric solutes and their retention on polybutadiene-coated alumina 3
Kasai, M., see Noguchi, K. 145
——, Uchida, T. 327
Kenney, D., see Cheung, A. P. 119
Keresztes, P., see Papp, E. 157
Kikuta, C., see Mascher, H. 417
Kok, W. Th., see Engelsma, M. 201
Kraak, J. C., see Stegeman, G. 547
——, see Tock, P. P. H. 185
Kurganov, A.
——, Kuzmenko, O., Davankov, V. A., Eray, B., Unger, K. K. and Trüdinger, U.
Effect of polystyrene coating on pore, structural and chromatographic properties of silica packings 391
Kuzmenko, O., see Kurganov, A. 391
Lahtonen, K., see Örn, G. 627
Lee, H. K., see Li, S. F. Y. 245
Li, S. F. Y.
——, Lee, H. K. and Ong, C. P.
Optimization of mobile phase composition for high-performance liquid chromatographic separation by means of the overlapping resolution mapping scheme 245
Lucarelli, C., see Belliardo, F. 371
Lundahl, P. see Yang, Q. 379
Lunte, C. E., see Scott, D. O. 461
Lynch, R. J., see Schoenmakers, P. J. 169
Manoha, M., see Rouchouse, A., 601
Maris, F.
——, Hindriks, R., Vink, J., Peeters, A., Vanden Driessche, N. and Massart, L.
Validation of an expert system for the selection of initial high-performance liquid chromatographic conditions for the analysis of basic drugs 211
Marko-Varga, G.
——, Dominguez, E., Hahn-Hägerdal, B. and Gorton, L.
Selective post-column liquid chromatographic determination of sugars in spent sulphite liquor with two enzymatic electrochemical detectors in parallel 423
Marr, J. G. D.
——, Seaton, G. G. R., Clark, B. J. and Fell, A. F.
Multiple absorbance ratio correlation—a new approach for assessing peak purity in liquid chromatography 289
Mascher, H.
—— and Kikuta, C.
Determination of amoxicillin in plasma by high-performance liquid chromatography with fluorescence detection after on-line oxidation 417
Massart, L., see Maris, F. 211
Miyasaka, K., see Yamakawa, Y. 319
Nihira, M., see Hayashida, M. 133
Nilsson, I., see Gustafsson, S. 75
Nilsson, L. B.
Regulation of peak compression effects for substituted benzamides in reversed-phase liquid chromatography 253

Noguchi, K., see Uchida, T. 327
——, Yanagihara, Y., Kasai, M. and Hirata, N.
Chromatographic characteristics of nucleic acid components on vinyl alcohol copolymer gel columns 145
Novák, I., see Belliardo, F. 371
Örn, G.
——, Lahtonen, K. and Jalonen, H.
Direct high-performance liquid chromatographic separation of (+)- and (−)-medetomidine hydrochloride with an α_1-acid glycoprotein chiral column 627
Ohkubo, A., see Sakuma, I. 223
Ohtani, T., see Uchida, T. 327
Okuyama, T., see Yamakawa, Y. 319
Ong, C. P., see Li, S. F. Y. 245
Oostervink, R., see Stegeman, G. 547
Ośmiałowski, K., see Kaliszan, R. 3
Papp, E.
—— and Keresztes, P.
Retention behaviour of mono- and dicarboxylic acids, carbohydrates and alcohols in ion-exclusion chromatography 157
Peeters, A., see Maris, F. 211
——, see Schoenmakers, P. J. 169
Pfeffer, W. D.
—— and Yeung, E. S.
Indirect fluorometric detection in open-tubular capillary column chromatography 401
Pfleiderer, B.
——, Albert, K. and Bayer, E.
Investigations by ^{29}Si cross-polarization magic angle spinning NMR spectroscopy of reaction pathways of silica gel polyfunctional modification 343
Poppe, H.
Secondary equilibria and their interaction with chromatographic transport 45
——, see Stegeman, G. 547
——, see Tock, P. P. H. 185
Quintero, G., see Vigh, Gy. 481
Rhodes, G. R., see Boppana, V. K. 279
Rouchouse, A.
——, Manoha, M., Durand, A., Thenot, J. P.
Direct high-performance liquid chromatographic determination of the enantiomers of alfuzosin in plasma on a second-generation α_1-acid glycoprotein chiral stationary phase 601
Sági, J., see Valkó, K. 35
Sakuma, I.
——, Takai, N., Dohi, T., Fukui, Y. and Ohkubo, A.
Resolution of unresolved peaks containing unknown components by high-performance liquid chromatography with multi-wavelength detection 223
Salvadori, P., see Betucci, C. 617
Schill, G., see Arvidsson, E. 579
Schoenmakers, P. J.
——, Peeters, A. and Lynch, R. J.
Optimization of chromatographic methods by a combination of optimization software and expert systems 169
——, Uunk, L. G. M. and Janssen, H.-G.
Comparison of stationary phases for packed-column supercritical fluid chromatography 563
Scott, D. O.
——, Sorensen, L. R. and Lunte, C. E.
In vivo microdialysis sampling coupled to liquid chromatography for the study of acetaminophen metabolism 461
Seaton, G. G. R., see Marr, J. G. D. 289
Siwek, A.
—— and Śliwiok, J.
Reversed-phase high-performance liquid chromatography in the investigation of the hydrophobicity of selected ketones 109
Śliwiok, J., see Siwek, A. 109
Smit, H. C., see Engelsma, M. 201
Sokolowski, A., see Fornstedt, T. 61
Sorensen, L. R., see Scott, D. O. 461
Ståhlberg, J., see Bartha, A. 85
Stegeman, G.
——, Oostervink, R., Kraak, J. C., Poppe, H. and Unger, K. K.
Hydrodynamic chromatography of macromolecules on small spherical non-porous silica particles 547
Szemző, A., see Valkó, K. 35
Takagi, T.
Application of low-angle laser light scattering detection in the field of biochemistry. Review of recent progress 409
Takai, N., see Sakuma, I. 223
Thenot, J. P., see Rouchouse, A. 601
Thompson, D. H., see Camacho, P. L. 611
Tock, P. P. H.
——, Duijsters, P. P. E., Kraak, J. C. and Poppe, H.
Theoretical optimization of open-tubular columns for liquid chromatography with respect to mass loadability 185
Trüdinger, U., see Kurganov, A. 391
Ucello-Barretta, G., see Bertucci, C. 617
Uchida, T.
——, Ohtani, T., Kasai, M., Yanagihara, Y., Noguchi, K., Izu, H. and Hara, S.
Peptide behaviour and analysis on a chemically stable C_{18}-bonded vinyl alcohol copolymer column with alkaline and acidic eluents 327
Unger, K. K., see Kurganov, A. 391
——, see Stegeman, G. 547

Uunk, L. G. M., see Schoenmakers, P. J. 563
Valkó, K.
——, Cserháti, T., Fellegvári, I., Sági, J. and Szemző, A.
Application of chromatographic retention data in an investigation of a quantitative structure–nucleotide incorporation rate relationship 35
Vanden Driessche, N., see Maris, F. 211
Vigh, Gy., see Bartha, A. 85
——, see Camacho, P. L. 611
——, Quintero, G. and Farkas, Gy.
Displacement chromatography on cyclodextrin–silicas. III. Enantiomer separations 481
——, see Welsch, T. 97
Vink, J., see Maris, F. 211
Wallstén, M., see Yang, Q. 379
Watanabe, T., see Hayashida, M. 133
Webster, R., see Camacho, P. L. 611
Welsch, T.
——, Frank, H. and Vigh, Gy.
Silanol effects in reversed-phase liquid chromatography 97
Westerlund, D., see Fornstedt, T. 61
Yamada, Y., see Yamakawa, Y. 319
Yamakawa, Y.
——, Miyasaka, K., Ishikawa, T., Yamada, Y. and Okuyama, T.
High-performance liquid chromatography of transfer ribonucleic acids on spherical hydroxyapatite beads 319
Yanagihara, Y., see Noguchi, K. 145
——, see Uchida, T. 327
Yang, Q.
——, Wallstén, M. and Lundahl, P.
Lipid-vesicle-surface chromatography 379
Yeung, E. S., see Pfeffer, W. D. 401